# The Roots of Anxiety and Panic



| THE ROOTS OF ANXIETY AND PANIC                       | 1        |
|--|----------|
| Introduction.  |          |
| Let's define it                                      |          |
| IF YOU HAVE ANXIETY AND/OR PANIC, YOU ARE NOT ALONE. |          |
| Underlying biology.                                  | 4        |
| Underlying psychology                                |          |
| Attentional Focus                                    |          |
| Cognitive beliefs                                    |          |
| What would predispose someone to anxiety or panic?   | <u>7</u> |
| References.  | 9        |
| RECOMMENDED READING.                                 | 10       |

## Introduction

You see a rattlesnake.

The image is sent to a part of your brain called the amygdala. It is the part of your brain that is primitive and thinks literally. When unregulated, it produces immediate and uncontrolled reactions. The hippocampus and the cortex, on the other hand, are able to process a stimulus *in context* by using memories of previous experiences. They allow us to react differently when we see a rattlesnake that is safely behind glass and one that is coiled at the base of our feet. Put another way, your amygdala is quick to put the gas pedal to the floor (raising your heart rate, speeding up your breathing), while the hippocampus can hit the brakes to help calm you down (Wallin, 2007, p. 72).

This passage from the book "Fear itself" describes the process in greater detail:

The primitive fear system kicks in around the first tenth of a second after our initial perception, prior to conscious awareness. The rational fear system begins to operate when sensory data processed by the cortex enter conscious awareness a fraction of a second later. [···] When we are in a dangerous situation, the primitive fear system stimulates an intense form of alertness to any potential threat.

 $[\cdots]$ 

Panic is the hallmark of the primitive fear system. This system evolved to meet the primal needs of the savage, brutish, life-or-death environment of one hundred thousand years ago, an environment in which most human beings never reached their thirtieth birthday. (Dozier, 1998, pp. 10-13)

# Let's define it

# Anxiety is:

- A state of unease, apprehension and worry.
- Disruptive to your life.
- May include physical symptoms such as heart palpitations, trembling and dry mouth

## Panic is:

- Intense fear
- You dread the possibility of another attack
- Symptoms can last for hours. Symptoms can include:

Difficulty breathing or feeling "smothered"

• Fear that you are "going crazy"

Racing heart or chest pain

Nausea or heartburn

• Feeling lightheaded or "not there"

Sweating

If you have anxiety and/or panic, you are not alone

According the 1999 report by the Surgeon General, anxiety disorders affect

about 16% of adults age 18-54. That's about 1 in every 6 people. Some type of panic

disorder affects about 1.5% of that same group. (U.S. Department of Health and

Human Services).

Having anxiety or panic doesn't mean that there is something wrong with you

or that you are "going crazy." To the contrary, it is a strong indicator from your

subconscious that something scary and dangerous has happened in your past and to

be careful in the future. And since phobias are often the result of a one time

experience, the fact that you remembered means that you are a fast learner (Andreas,

Faulkner, 1994, p. 194). Learning quickly has a strong survival value. Remember, the

zebra that doesn't learn to check for crocodiles before drinking at the river doesn't

live very long.

Michael Etts, LCSW-C, ACH www.AdaptiveTherapy.com Underlying biology

There are several medical issues that could be causing or contributing to

anxiety and panic. These are hypoglycemia, hyperthyroidism, mitral valve prolapse,

PMS and inner ear disturbances, to name a few (Bourne, 2005, pp.43-44).

The biological causes of anxiety and panic are the subject of a growing body of

evidence. I will note two critical structures that are implicated in anxiety and panic

(see the <u>recommended reading</u> for additional information). A key structure in the

brain is the hippocampus. When adrenal steroids (stress hormones) reach the

hippocampus, it starts a chain reaction that eventually inhibits additional adrenal

activity. Hence, the hippocampus is important in modulating stress response. Current

research indicates that prolonged periods of stress, such as those experienced in child

abuse or neglect, cause damage to the hippocampus. Vietnam veterans and survivors

of long term child abuse actually have a shrunken hippocampus, as well as impaired

memory (LeDoux, 1996, pp. 240-242). But this damage appears to be at least

somewhat reversible, as recent research shows that the hippocampus can regenerate

nerve cells (Bremmer, Narayan, 1998).

In contrast, the positive effects of "maternal care, handling and soothing touch

[...] protect the hippocampus from stress" (Cozolino, 2006, p. 236). For more

Michael Etts, LCSW-C, ACH

information on the effects of early maternal care and stress, see the April 2008 newsletter <u>Early childhood and the ability to cope with trauma</u>.

A second brain structure related to fear and anxiety is the amygdala. As stated previously, the amygdala gets sensory input first and, if is a hint of a threat, it begins to prime your body for reaction. Research suggests that the amygdala has an indelible memory. Once a major stressful or traumatic event has occurred, the amygdala records it forever (LeDoux, 1996). While this might seem like a hopeless obstacle to recovery, it can be counterbalanced by new learning. For example, imagine someone is bitten by a dog and subsequently becomes afraid of dogs. Over time, if that person begins approaching friendly dogs and they can create new, more positive memories. These memories are laid down in the brain as new knowledge, such as "most dogs don't bite". This new learned knowledge can then act as a counterbalance to the amygdala's initial response of fear. There are several psychological techniques (to be discussed in the next newsletter, the Treatment of Anxiety and Panic) that can facilitate this new learning.

While both the hippocampus and the amygdala are affected by stress and trauma, they appear to be affected in opposite ways. The hippocampus will be less likely to record a clear memory of the event, while the amygdala is apt to

your fear center may remember a trauma that you cannot...

Michael Etts, LCSW-C, ACH
 www.AdaptiveTherapy.com
 2008 Adaptive Therapy, All rights reserved

lay down a strong and permanent record. Again, from

LeDoux "It is thus completely possible that one might have

poor conscious memory of a traumatic experience, but at

the same time form a very powerful implicit, unconscious

emotional memories [...]" and "[...] if a conscious memory

wasn't formed, it can't be recovered" (pp. 245-246). In

other words, your fear center may remember a trauma that

you cannot.

Underlying psychology

There are many psychological features that are associated with persistent

anxiety and panic. Two major categories are attentional focus and cognitive beliefs.

Attentional focus is what a person directs their attention to. Cognitive beliefs guide

your interpretation of events by a schema or belief system.

Attentional Focus

There is far too much information coming in from all our senses to be

perceived all at once. At any moment, you are focused on some aspects of the reality

in front of you and selectively ignoring other parts. From "On Intelligence":

Nichael Etts, LCSW-C, ACH

At any moment in time, you can directly sense only a tiny part of your world.

That tiny part dictates what memories will be invoked, but it isn't sufficient on

its own to build the whole of your current perception. […]

It is my memory model of the world that predicts [...] by analogy to past

experience. Most of what you perceive is not coming through your senses; it is

generated by your internal memory model. (Hawkins, p. 202)

If someone is scanning the social or physical environment for danger cues, they

will be downplaying or missing safety cues. Take a bookshelf with paperweights on it.

The anxious person might see the paperweights as potential blunt force weapons

while a nonanxious might focus on the book titles.

Attentional focus can also be internal. Being keenly aware of bodily sensations

and interpreting those physical sensations negatively is thought make someone more

vulnerable to panic attacks (Barlow, 2002).

Internal and external attentional focus can be illustrated by someone who is

anxious about public speaking. Externally, the person might locate a few people who

look uninterested or bored and then lose sight of the part of the audience that is

showing interest. Internally, the person notices tightness in their stomach and begins

to focus on it, ignoring the relaxation that they feel in other parts of their body.

Michael Etts, LCSW-C, ACH

- 8 -

Negative interpretations of these two focuses will increase the felt sense of anxiety and the possibility of panic. So it is not just what we perceive but also how we interpret those perceptions.

# Cognitive beliefs

Anxiety is supported by a view that the world is a dangerous place, and having that view direct your attention toward threat and away from evidence of safety. Combine this danger focus with the belief that you will be unable to cope with the threatened harm and you have a case of escalating anxiety. As the anxiety rises, it begins to create a kind of tunnel vision, through which the ability to reason and assess the situation rationally is degraded. This heightened state makes it even more difficult to see external safety cues or to recall comforting and reassuring memories. Now, add in any of the symptoms of panic (light headedness, heart racing, nausea, sweating) and the likely interpretation would be that a panic attack is arriving.

An excerpt from the study "Attentional Fixation in Panic Disorder" states it succinctly:

Anxious individuals are characterized by the heightened activation of a "danger" schema, which orients them toward threat in their environment and directs cognitive resources toward processing threat cues at the expense of safety cues. Schemas also comprise beliefs about the world being dangerous and about the inability to cope with potential harm. When activated, these beliefs

Michael Etts, LCSW-C, ACH www.AdaptiveTherapy.com prompt catastrophic misinterpretations that occur in uncomfortable situations

(Wenzel, Sharp, Sokol, Beck, 2006, p.65).

What would predispose someone to anxiety or panic?

Three major factors that would predispose someone to anxiety or panic are

social learning, stress and insecure attachment.

Social learning would include factors like witnessing a heart attack or cancer in

a close relative and then becoming anxious about developing that condition. Studies

also show that anxious adults are more likely to have had anxious caregivers or

caregivers who encourage sick role behavior. (Barlow, 2002, p. 225).

Early stress is another major factor related to anxiety and panic. Barlow writes

in "Anxiety and its disorders":

Now, there is dramatic evidence that early stressful life events may effect rather

permanent alterations in brain function [...] that may mediate the

neurobiological vulnerability to develop chronic anxiety and depression later in

life (p. 217).

As discussed in the biology section, early traumatic experience can actually alter the

structure of the brain.

Michael Etts, LCSW-C, ACH

A person's attachment type (secure vs. insecure, see the April 2008 newsletter

for more details) is critical in determining what resources they will have available

when confronting stressful life situations. In insecure/disorganized attachment, it

appears that a caregivers frighten their children not only with frightening behavior,

but also with frightened behavior. In situations where the caregiver's own trauma or

loss is triggered, they can behave in ways that transmit that fright. It is theorized that

this may be how can intergenerational transmission of fearfulness occurs. (Wallin,

2007).

In the next newsletter, I will address the treatment of anxiety and panic.

Michael Etts, LCSW-C, ACH www.AdaptiveTherapy.com

# References

Andreas, S. Faulkner, C. (1994). *NLP: The New Technology of Achievement*. New York: HarperCollins Publishing.

Barlow, D. (2002). *Anxiety and its disorders: The nature and treatment of anxiety and panic* (2nd ed.). New York: Guilford Press.

Bourne, E. (2005). *The Anxiety & Phobia Workbook*. California: New Harbinger Publications.

Bremmer, J. D., Narayan M. (1998). The Invisible Epidemic: Post-Traumatic Stress Disorder, Memory and the Brain. *Developmental Psychopath*ology. 10:871-885.

Cozolino, L. (2006). *The Neuroscience of Human Relationships: Attachment and the Developing Social Brain.* New York: W.W. Norton & Company.

Dozier, R. (1998). Fear Itself: The Origin and Nature of the Powerful Emotion that Shapes Our Lives and Our World. New York: St Martins Press.

Hawkins, J. (2005) On Intelligence. New York: Owl Books/Henry Holt & Co.

Jemmer, P. (2005). Phobia: Fear and Loathing in Mental Spaces. *European Journal of Clinical Hypnosis*, 6(3), 24–32.

LeDoux, J. (1996). *The Emotional Brain: The Mysterious Underpinnings of Emotional Life.* New York: Simon & Schuster.

Satcher, D. (1999). *Mental Health: A Report of the Surgeon General* Rockville, MD: U.S. Department of Health and Human Services,

Substance Abuse and Mental Health Services Administration, Center for Mental Health Services, National Institutes of Health,

National Institute of Mental Health, 1999, Washington, DC. Retrieved July 14, 2008 from <a href="http://www.surgeongeneral.gov/library/reports.htm">http://www.surgeongeneral.gov/library/reports.htm</a>

Wallin, D. (2007) Attachment in Psychotherapy. New York: The Guilford Press.

Wenzel A, Sharp I, Sokol L, Beck A. (2006) Attentional Fixation in Panic Disorder. *Cognitive Behaviour Therapy.* 35(2), pp. 65 – 73.

# Recommended reading

Self help for anxiety and panic

The Anxiety & Phobia Workbook

A self help encyclopedia for anxiety, phobias and panic

For those seeking a deeper understanding and the research underpinnings:

Biological aspects of anxiety and panic:

The Emotional Brain: The Mysterious Underpinnings of Emotional Life.

Written by a brain researcher, this book examines the brain structures that are associated with fear and anxiety.

For an exhaustive overview of anxiety

Anxiety and Its Disorders: The Nature and Treatment of Anxiety and Panic

A tome of research and theory with extensive information about anxiety and panic

## Services

# More information

Anyone who has ever had to deal with weeds knows that it does no good to keep cutting off the tops. You have to get to the *root*. It's the same concept when you are dealing with psychological problems.

If you are ready to begin removing the legacy of trauma, if you are ready to start making anxiety, depression and insomnia a part of your past, then make a call and schedule a consultation.

© Michael Etts, 2008. All Rights Reserved.

Michael Etts, LCSW-C, ACH

On the web: <a href="http://www.adaptivetherapy.com">http://www.adaptivetherapy.com</a>

Email: <u>mike@adaptivetherapy.com</u>

Phone: 1-5240-253-7051

To URL of this newsletter, click on the link below:

http://www.adaptivetherapy.com/The%20roots%20of%20anxiety%20and %20panic.pdf